

George Quintero, D.D.S.

Diplomat, American Board of Periodontology
Diplomat, American Board of Oral Medicine
Fellow, Academy of General Dentistry

Scaling and Root Planing

You have had Scaling and Root Planing in your mouth and you can expect normal healing events over the next week or two. The roots affected by disease have been thoroughly cleaned and are no longer covered by diseased gum tissue. Because of this, you will probably experience temperature sensitivity. This is all normal and temporary.

You may have been given a prescription for a strong pain medication, or an over-the-counter pain medication such as Tylenol or Ibuprofen. This will allow you to sleep at night and function fairly normally during the day. Do not drive or operate machinery when you take this medication as you will be drowsy. To avoid nausea, have some food in your stomach when you take this medication. Do not take any more of your medication than you need in order to control pain, and take it as directed. We have done everything we can to minimize pain.

Expect some bleeding during the first 12 hours. If you rest and take it easy, this should be minimal. If the bleeding is coming from one specific point, you may apply pressure directly over the area with some damp gauze.

A soft diet for two or three days is recommended. Please do not eat peanuts, popcorn, wild rice, or any foods with shells or husks.

Begin the oral hygiene procedure the night of Scaling and Root Planing. Run hot water over your bristles to soften them before gently brushing the treated area. The cleaner you keep your teeth and gums during the first week, the better they will feel and heal. Expect a little bleeding when you begin to clean your teeth and gums. This is normal and you are not doing any harm. You may want to use a salt water rinse 3 times a day during the first week. This is prepared by mixing a teaspoon of salt in a medium size (8 oz.) glass of lukewarm water.

TIPS

When you go home take your prescribed pain medication, or an over-the-counter pain medication such as Tylenol or Ibuprofen with food and water before anesthesia wears off

Soft diet for 2 to 3 days

Cold foods such as ice cream, applesauce, yogurt, pudding are good things to eat when you get home

NO SMOKING or carbonated beverages for 2 days

No peanuts, popcorn, wild rice or foods with shells, husks or spicy food

Take medication with food and water

Questions call George Quintero, D.D.S. Office 770-614-8823 Cell phone 770-318-0703

Pain Medication

You may have been given a prescription for a very strong pain medication. I have prescribed a generic drug since it works as well as the brand name equivalent and costs considerably less. Be sure the pharmacist gives you the generic brand. If a refill has been requested on your prescription, be sure the pharmacist notes this on the label.

Be sure to take your medication as directed. An 800mg tablet of Ibuprofen or four (200mg) tablets of Ibuprofen can be taken with or in place of the pain medication. The Ibuprofen provides additional pain relief. Be sure to have food in your stomach before taking either medication.

TIPS

Take one (1) tablet of the prescribed pain medication for your first dose until you are certain how this medication affects you. Take only as much medication as necessary to control your discomfort. Sometimes the relief only lasts 2-3 hours. Taking Ibuprofen with the prescribed medication will help prolong the relief.

If this medication makes you nauseous, take 1/2 tablet with food in your stomach followed 30 minutes later by another 1/2 tablet. This allows the medication to enter your body slowly.

There are some other common side-effects that you may experience, if you take more of the pain medication than your body can effectively handle. You may experience sweating, headaches, and insomnia. If you are experiencing any of these annoying side-effects, you can eliminate them by reducing the amount of pain medication you are taking. *If you experience any signs of an allergic reaction, stop taking the medication and call Dr. Quintero.*

Do not take any more medications than you need to manage your discomfort. It will only increase side-effects. We can not prescribe an unlimited supply of this medication.

Questions call George Quintero, D.D.S. Office 770-614-8823 Cell phone 770-318-0703

Email: drq@quinteroperio.com • www.quinteroperio.com